

GANEA AND WHITEHEAD DEFINITIONS FOR THE TANGENTIAL LUSTERNIK-SCHNIRELMANN CATEGORY OF FOLIATIONS

JEAN-PAUL DOERAENE, ENRIQUE MACIAS-VIRGÓS, AND DANIEL TANRÉ

ABSTRACT. This work solves the problem of elaborating Ganea and Whitehead definitions for the tangential category of a foliated manifold. We develop these two notions in the category $\mathcal{S}\text{-Top}$ of stratified spaces, that are topological spaces X endowed with a partition \mathcal{F} and compare them to a third invariant defined by using open sets. More precisely, these definitions apply to an element (X, \mathcal{F}) of $\mathcal{S}\text{-Top}$ together with a class \mathcal{A} of subsets of X ; they are similar to invariants introduced by M. Clapp and D. Puppe.

If $(X, \mathcal{F}) \in \mathcal{S}\text{-Top}$, we define a transverse subset as a subspace A of X such that the intersection $S \cap A$ is at most countable for any $S \in \mathcal{F}$. Then we define the Whitehead and Ganea LS-categories of the stratified space by taking the infimum along the transverse subsets. When we have a closed manifold, endowed with a C^1 -foliation, the three previous definitions, with \mathcal{A} the class of transverse subsets, coincide with the tangential category and are homotopical invariants.

CONTENTS

1. The category of stratified spaces	3
2. Lusternik-Schnirelmann category of stratified pairs	6
3. Lusternik-Schnirelmann category of stratified spaces	10
4. Tangential LS-category of foliations	11
References	13

2000 *Mathematics Subject Classification.* 55M30-57R30-55U35.

Key words and phrases. LS-category. Closed model category. Foliation. Tangential category. Stratified space.

The second author is partially supported by research project MICINN MTM2008-05861.

The Lusternik-Schnirelmann category (LS-category in short) of a smooth manifold M is an invariant putting in relation the topological complexity of M with the behavior of smooth functions defined on M . In particular, when the manifold is compact, it is a lower bound of the number of critical points for any smooth function. This numerical invariant can also be defined for a topological space X and reveals itself as a homotopy type invariant. The LS-category $\text{cat}(X)$ is the least natural number n such that there exists a cover of X by $n + 1$ open sets, each of them being contractible to a point inside X .

H. Colman and the second author ([5] and [6], see also [3]) have adapted this definition to the case of a foliated manifold (M, \mathcal{F}) . Indeed, they introduced two invariants, one which refers to the transverse structure and a second one, the *tangential category*, $\text{cat}_{\mathcal{F}}(M)$, see Definition 18 for a precise statement. Our paper is concerned with this tangential category, also investigated by W. Singhof and E. Vogt [15]. These two authors prove that the tangential category is less than or equal to $\dim \mathcal{F} + 1$, a result similar to the fact that the dimension of a CW-complex plus 1 is an upper bound for its LS-category. Some other results of [15] will be crucial for our work and we will quote them precisely in the last section. In [4], H. Colman and S. Hurder prove, for instance, that the nilpotency index of the reduced filtered cohomology is a lower bound of the tangential category.

In [11], S. Hurder says that the homotopy-theoretic interpretation of $\text{cat}_{\mathcal{F}}(M)$ corresponding to the Whitehead and Ganea definitions of category is “one of the most important open problems in the subject”. In the present work we give for the tangential category several equivalent definitions inspired by the Whitehead procedure (using the fat wedge), and by the Ganea construction (using Milnor’s classifying spaces). As in the classical case, we do that in the topological framework, considering the most general setting of topological spaces, endowed with a partition. We call them *stratified spaces* and denote by **S-Top** the category of stratified spaces and stratified maps.

Before giving the content of the paper, we present our general strategy. Recall that the LS-category of a space X uses open sets that are contractible to a point inside X . If X is path-connected, one can use contractions on a specific point and work with pointed tools as fat wedges and Ganea fibrations. These types of constructions are also present in our work and, at a first look, we should distinguish some transverse structure for a foliation. But we cannot do this, because the definition of tangential LS-category is made of tangentially contractible open sets without specific transverse set. We solve this contradiction by taking an adequate definition of topological transverse set and the infimum on all transverse sets. It is a priori unclear that this process gives the tangential LS-category of a foliation and Section 4 contains a proof of that fact.

The second main point is the equality between the invariants coming from the Whitehead and Ganea constructions. As the first author showed in [8], one has only to prove that **S-Top** is a closed model category satisfying the Cube Lemma. Instead of that, we take here a shortcut: the heart of the

Cube Lemma needs only a structure of fibration category as it appears in the proof of Theorem B. Therefore, in Theorem A of Section 1, we prove that $\mathcal{S}\text{-Top}$ is a category of fibrations, which is sufficient for our purpose.

In Section 2, we consider a stratified pair (X, A, \mathcal{F}_X) , where (X, \mathcal{F}_X) is a stratified space and A a subset of X . We introduce the Whitehead and Ganea constructions which give the notions of Whitehead and Ganea category, respectively denoted $\text{Whcat}(X, A, \mathcal{F}_X)$ and $\text{Gcat}(X, A, \mathcal{F}_X)$. We prove their equality in Theorem B. These constructions are adaptations of those by M. Clapp and D. Puppe [2] in the case of topological spaces. In a third step, we want a notion of LS-category for (X, A, \mathcal{F}_X) using open sets. An open subset U of X is said *A-categorical* if there is a stratified homotopy defined on (U, \mathcal{F}_U) between the identity on U and a map with values in A , see Definition 7. (The induced stratification \mathcal{F}_U is composed of the connected components of the intersections $U \cap S$, $S \in \mathcal{F}_X$.) This notion of *A-categorical* open set brings a definition of open LS-category, $\text{Ocat}(X, A, \mathcal{F}_X)$, in the usual way. When A is a *B* stratified neighborhood deformation (in short *B-SND*, see Definition 11) and X a normal space, we prove that $\text{Ocat}(X, B, \mathcal{F}_X) \leq \text{Whcat}(X, A, \mathcal{F}_X) \leq \text{Ocat}(X, A, \mathcal{F}_X)$, see Theorem C.

In Section 3, we introduce the *transverse subsets* which are the key for the comparison with the tangential LS-category of foliations. If $(X, \mathcal{F}_X) \in \mathcal{S}\text{-Top}$, a subset A of X is called *transverse to \mathcal{F}_X* if the intersection $A \cap S$ is at most countable for all strata $S \in \mathcal{F}_X$. The transverse subsets to a foliation own this property, see [15]. In the definition of the tangential LS-category of foliations, there is no predetermined transverse set A , so we define $\text{Ocat}(X, \mathcal{F}_X)$ as the infimum of the integers $\text{Ocat}(X, A, \mathcal{F}_X)$ when A is transverse to \mathcal{F}_X (analogously for $\text{Whcat}(X, A, \mathcal{F}_X)$ and $\text{Gcat}(X, A, \mathcal{F}_X)$). In Theorem D, we prove that $\text{Ocat}(X, \mathcal{F}_X)$ is a homotopy invariant in $\mathcal{S}\text{-Top}$.

Finally, in Section 4, we consider a smooth closed manifold with a C^1 -foliation, (M, \mathcal{F}_M) , and prove the equalities:

$$\text{cat}_{\mathcal{F}}(M) = \text{Ocat}(M, \mathcal{F}_M) = \text{Gcat}(M, \mathcal{F}_M) = \text{Whcat}(M, \mathcal{F}_M),$$

see Theorem E. We end with the example of the Reeb foliation on the torus T^2 by giving a transverse set A that is an *A-SND*.

The second author would like to thank the University of Sciences and Technology of Lille and the first and third authors the University of Santiago de Compostela for their hospitality during the realization of this work.

1. THE CATEGORY OF STRATIFIED SPACES

A *stratified space* (X, \mathcal{F}_X) is a topological space together with a partition \mathcal{F}_X whose elements $S \subseteq X$ are path-connected subspaces. We denote by $\sim_{\mathcal{F}_X}$ (or \sim_X if there is no ambiguity) the equivalence relation associated to the partition \mathcal{F}_X of X and call *strata* the elements S of \mathcal{F}_X . A stratified map between stratified spaces is a continuous map compatible with the equivalence relations.

Let **Top** be the category of topological spaces and continuous maps and **S-Top** be the category of stratified spaces and stratified maps. We first observe that the category **S-Top** has finite direct and inverse limits. This is obvious from the corresponding limits in **Top**.

The stratification on a pull-back $(X, \mathcal{F}_X) \times_{(B, \mathcal{F}_B)} (Y, \mathcal{F}_Y) = (P, \mathcal{F}_P)$ is defined by $(x, y) \sim_P (x', y')$ if, and only if,

$$x \sim_X x' \text{ and } y \sim_Y y'.$$

If we have a commutative diagram in **S-Top**:

$$\begin{array}{ccc} (Z, \mathcal{F}_Z) & \xrightarrow{j} & (Y, \mathcal{F}_Y) \\ \downarrow i & & \downarrow g \\ (X, \mathcal{F}_X) & \xrightarrow{f} & (B, \mathcal{F}_B) \end{array}$$

then the map induced by the pull-back in **Top**,

$$(i, j) : Z \rightarrow P = X \times_B Y,$$

is a stratified map in **S-Top**, because if $z \sim_Z z'$, then $i(z) \sim_X i(z')$ and $j(z) \sim_Y j(z')$, so $(i, j)(z) = (i(z), j(z)) \sim_P (i(z'), j(z')) = (i, j)(z')$.

The stratification on a push-out $(X, \mathcal{F}_X) \vee_{(B, \mathcal{F}_B)} (Y, \mathcal{F}_Y) = (S, \mathcal{F}_S)$ of $f : (B, \mathcal{F}_B) \rightarrow (X, \mathcal{F}_X)$ and $g : (B, \mathcal{F}_B) \rightarrow (Y, \mathcal{F}_Y)$ is defined by

- $x \sim_S x'$ if, and only if,
 - either: $x \sim_X x'$,
 - or: $x \sim_X f(b)$ and $x' \sim_X f(b')$ with $g(b) \sim_Y g(b')$, (†)
- a similar formula for $y \sim_S y'$,
- and $x \sim_S y$ if, and only if,
 - $x \sim_X f(b)$ and $y \sim_Y g(b)$ (‡).

If we have a commutative diagram in **S-Top**:

$$\begin{array}{ccc} (B, \mathcal{F}_B) & \xrightarrow{g} & (Y, \mathcal{F}_Y) \\ \downarrow f & & \downarrow j \\ (X, \mathcal{F}_X) & \xrightarrow{i} & (Z, \mathcal{F}_Z) \end{array}$$

then the map induced by the push-out in **Top**,

$$[i, j] : S = X \vee_B Y \rightarrow Z,$$

is a stratified map in **S-Top**, because if $x \sim_S x'$, with condition (†), then $i(x) \sim if(b) = jg(b) \sim jg(b') = if(b') \sim i(x')$ and if $x \sim_S y$, with condition (‡), then $i(x) \sim if(b) = jg(b) \sim j(y)$.

Denote by I the interval $[0, 1]$ endowed with the trivial partition $\mathcal{F}_I = \{[0, 1]\}$ and define a notion of homotopy in **S-Top** as follows.

Definition 1. Two stratified maps, $f, g : (X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$, are called **S-homotopic** if there exists a stratified map $H : (X, \mathcal{F}_X) \times (I, \mathcal{F}_I) \rightarrow (Y, \mathcal{F}_Y)$ such that $f = H(-, 0)$ and $g = H(-, 1)$. We denote this relation by $f \simeq_S g$.

Recall that **Top** can be endowed with two structures of closed model category. The first one (see [14]) has for fibrations the Serre fibrations and for weak equivalences the classical weak equivalences. The second one (see

[16]) has for fibrations the Hurewicz fibrations and for weak equivalences the homotopy equivalences. For the study of $\mathcal{S}\text{-Top}$, we consider a situation “à la Hurewicz” and prove that $\mathcal{S}\text{-Top}$ is a fibration category, in the sense of Baues (see [1, I §1a]), which is sufficient for our objective.

Theorem A. *The category $\mathcal{S}\text{-Top}$ is a fibration category with*

- fibrations, *the stratified maps $f: (X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$ having the \mathcal{S} -homotopy lifting property;*
- weak equivalences, *the stratified maps $f: (X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$ such that there exists a stratified map $g: (Y, \mathcal{F}_Y) \rightarrow (X, \mathcal{F}_X)$ with $f \circ g \simeq_{\mathcal{S}} \text{id}_Y$ and $g \circ f \simeq_{\mathcal{S}} \text{id}_X$.*

Note that the full subcategory of stratified spaces (X, \mathcal{F}_X) with \mathcal{F}_X formed of the path-connected components of X equals \mathbf{Top} with its structure of fibration category. In particular, a weak equivalence in $\mathcal{S}\text{-Top}$ is a weak equivalence in \mathbf{Top} .

Proof. As the proof is an adaptation of [16] to the stratified case, we only sketch it. Recall that a fibration category is a category satisfying the axioms (F1), (F2), (F3) and (F4) of [13].

(F1) The isomorphisms are weak equivalences and also fibrations. Given two maps

$$A \xrightarrow{f} B \xrightarrow{g} C$$

if any two of f, g, gf are weak equivalences, then so is the third. The composite of fibrations is a fibration. The verification of these properties is immediate.

(F2) Consider a pull-back in $\mathcal{S}\text{-Top}$:

$$\begin{array}{ccc} (P, \mathcal{F}_P) & \xrightarrow{g} & (E, \mathcal{F}_E) \\ q \downarrow & & \downarrow p \\ (Y, \mathcal{F}_Y) & \xrightarrow{f} & (B, \mathcal{F}_B) \end{array}$$

We have to show that if p is a fibration of $\mathcal{S}\text{-Top}$, then so is q . Indeed, from the universal property, the map q has the \mathcal{S} -homotopy lifting property and therefore is a fibration of $\mathcal{S}\text{-Top}$.

We have also to show that if p is a trivial fibration of $\mathcal{S}\text{-Top}$, then so is q . As p is a weak equivalence, there exists $\sigma: (B, \mathcal{F}_B) \rightarrow (E, \mathcal{F}_E)$ such that $p \circ \sigma \simeq_{\mathcal{S}} \text{id}_B$ and $\sigma \circ p \simeq_{\mathcal{S}} \text{id}_E$. As p is a fibration, we may suppose that $p \circ \sigma = \text{id}_B$. From the universal property, we deduce the existence of $\mu: (Y, \mathcal{F}_Y) \rightarrow (P, \mathcal{F}_P)$ such that $g \circ \mu = \sigma \circ f$ and $q \circ \mu = \text{id}_Y$. We have to show that $\mu \circ q \simeq_{\mathcal{S}} \text{id}_P$. For that, we follow the proof in the case of the category \mathbf{Top} [16], adding the ad hoc argument for the stratification. We briefly recall the main steps for the convenience of the reader:

- First, one knows (see [9]) that the homotopy H between $\sigma \circ p$ and the identity on E can be chosen such that $p \circ H(-, t) = p$ for any $t \in I$.

- With that homotopy H , we check the commutativity of the following diagram, where proj means a projection onto one factor.

$$\begin{array}{ccccc}
 & & P \times I & \xrightarrow{g \times \text{id}} & E \times I \\
 & & \downarrow & & \downarrow \\
 & & P & \xrightarrow{g} & E \\
 & & \downarrow q \times \text{id} & & \downarrow p \times \text{id} \\
 & & Y \times I & \xrightarrow{f \times \text{id}} & B \times I \\
 & & \downarrow \text{proj} & & \downarrow \text{proj} \\
 P & \xrightarrow{q} & Y & \xrightarrow{f} & B
 \end{array}$$

From the universal property, there exists $G: (P, \mathcal{F}_P) \times (I, \mathcal{F}_I) \rightarrow (P, \mathcal{F}_P)$ such that $g \circ G = H \circ (g \times \text{id})$ and $q \circ G = \text{proj} \circ (q \times \text{id})$. This map G satisfies $G(-, 1) = \text{id}$ and $G(-, 0) = \mu \circ q$ as wanted.

On the other hand, it is easily checked that any object is fibrant, which implies, as quoted by H. Baues (see [1, Lemma 1.4 p.7]), that the category $\mathcal{S}\text{-Top}$ is ‘proper’, i.e. if we assume that f is a weak equivalence, so is g .

(F3) Let $f: (X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$ be a map in $\mathcal{S}\text{-Top}$. We have to write this map as $f = p \circ s$ where s is a weak equivalence in $\mathcal{S}\text{-Top}$ and p is a fibration in $\mathcal{S}\text{-Top}$. Recall that Y^I denotes the space of continuous maps $\omega: I \rightarrow Y$. Here we denote by $(Y, \mathcal{F}_Y)^I$ the set of continuous maps, $\omega: I \rightarrow Y$, such that $\omega(I)$ is included in an element S of \mathcal{F}_Y , together with the partition formed of the subsets S^I with $S \in \mathcal{F}$. The proof of the axiom (F3) follows the classical way [16], substituting Y^I by $(Y, \mathcal{F}_Y)^I$. The different steps of the proof are:

- The map $p_1: (Y, \mathcal{F}_Y)^I \rightarrow (Y, \mathcal{F}_Y)$ defined by $p_1(\omega) = \omega(1)$ is a fibration in $\mathcal{S}\text{-Top}$.
- The map $s_1: (Y, \mathcal{F}_Y) \rightarrow (Y, \mathcal{F}_Y)^I$, sending y to the constant path \hat{y} , is an \mathcal{S} -homotopy inverse of p_1 .
- If we denote by P_f the pull-back of p_1 and f , the map $p_0: P_f \rightarrow Y$ defined by $p_0(\omega, x) = \omega(0)$ is a fibration.
- The map $s_f: (X, \mathcal{F}_X) \rightarrow P_f$, given by $x \mapsto (\widehat{f(x)}, x)$, is a weak equivalence of $\mathcal{S}\text{-Top}$ that gives the desired decomposition of f as $f = p_0 \circ s_f$.

(F4) We check easily that every trivial fibration of $\mathcal{S}\text{-Top}$ has a section, which implies (F4). \square

2. LUSTERNIK-SCHNIRELMANN CATEGORY OF STRATIFIED PAIRS

Definition 2. Let $(X, \mathcal{F}_X) \in \mathcal{S}\text{-Top}$ be a stratified space and A be a topological subspace of X . The *induced stratification* (A, \mathcal{F}_A) has for strata the connected components of the intersections $A \cap S$, with $S \in \mathcal{F}_X$. The pair (A, \mathcal{F}_A) is called a *stratified subspace* of (X, \mathcal{F}_X) and (X, A, \mathcal{F}_X) a *stratified pair*.

If $f: (X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$ is a stratified map, we observe from the previous definition, that the restriction $f: (X, \mathcal{F}_X) \rightarrow (f(X), \mathcal{F}_{f(X)})$ is also stratified.

2.1. Ganea construction: the invariant $\text{Gcat}(X, A, \mathcal{F})$.

Definition 3. The i th *Ganea space* $G_i(X, A)$ of a stratified pair (X, A, \mathcal{F}_X) is the set of $(i + 1)$ -uples $(\alpha_1, \dots, \alpha_{i+1})$ of paths in $(X, \mathcal{F}_X)^I$, such that $\alpha_1(1) = \dots = \alpha_{i+1}(1)$ and there exists k , $1 \leq k \leq i + 1$, with $\alpha_k(0) \in A$.

We observe from this definition that, for any element $(\alpha_1, \dots, \alpha_{i+1})$ of $G_i(X, A)$, there is an associated element S of the partition \mathcal{F}_X such that $\alpha_j(I) \subset S$, for all $j = 1, \dots, i + 1$.

We put on $G_i(X, A)$ the stratification induced by the product stratification and the stratification of X^I , and denote by $G_i(X, A, \mathcal{F}_X)$ this stratified space. The i th *Ganea fibration* $g_i: G_i(X, A, \mathcal{F}_X) \rightarrow (X, \mathcal{F}_X)$ is defined by $g_i(\alpha_1, \dots, \alpha_{i+1}) = \alpha_1(1)$.

Definition 4. The *Ganea LS-category* of a stratified pair (X, A, \mathcal{F}_X) is the least integer n such that the n th Ganea fibration $g_n: G_n(X, A, \mathcal{F}_X) \rightarrow (X, \mathcal{F}_X)$ has a stratified section. We denote it by $\text{Gcat}(X, A, \mathcal{F}_X)$.

2.2. Whitehead construction: the invariant $\text{Whcat}(X, A, \mathcal{F}_X)$.

Definition 5. The i th *fat wedge* of a stratified pair (X, A, \mathcal{F}_X) is the subset $T_i(X, A)$ of the product X^{i+1} formed of the elements (x_1, \dots, x_{i+1}) such that there exists k , $1 \leq k \leq i + 1$, with $x_k \in A$.

We endow it with the stratification induced from the stratification of the product $(X, \mathcal{F}_X)^{i+1}$ and denote this stratified space by $T_i(X, A, \mathcal{F}_X)$.

Definition 6. The *Whitehead LS-category* of a stratified pair (X, A, \mathcal{F}_X) is the least integer n such that the diagonal $\Delta: (X, \mathcal{F}_X) \rightarrow (X, \mathcal{F}_X)^{n+1}$ factors up to \mathcal{S} -homotopy through the inclusion map $t_n: T_n(X, A, \mathcal{F}_X) \rightarrow (X, \mathcal{F}_X)^{n+1}$. We denote it by $\text{Whcat}(X, A, \mathcal{F}_X)$.

If $A = \{*\}$, they coincide with the classical ones, see [7], [8], [2] or [10] for more details.

2.3. Open LS-category: the invariant $\text{Ocat}(X, A, \mathcal{F}_X)$. In this paragraph, we provide a third definition using open sets. First we precise the notion of categorical subsets in the context of stratified spaces.

Definition 7. Let (X, A, \mathcal{F}_X) be a stratified pair. A non-empty subset U of X is said to be *A-categorical* if there is a stratified homotopy $H: (U, \mathcal{F}_U) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)$ such that $H(x, 0) = x$ and $H(x, 1) \in A$ for any $x \in U$. We call H a *stratified deformation of U into A* .

Observe that, as in the classical case, the sets $H(U, t)$, with $t \in I$, are not necessarily contained in U . With the previous definition, we introduce the *open LS-category* of the stratified pair (X, A, \mathcal{F}_X) .

Definition 8. The *open LS-category* of a stratified pair (X, A, \mathcal{F}_X) is the least integer n such that there exists a covering of X by $(n + 1)$ open sets which are *A-categorical*. We denote it by $\text{Ocat}(X, A, \mathcal{F}_X)$.

We write also $\text{Ocat}(X, A, \mathcal{F}_X) = \infty$ if such a covering does not exist. Observe that $\text{Ocat}(X, X, \mathcal{F}_X) = 0$ and that the finiteness of $\text{Ocat}(X, A, \mathcal{F}_X)$ implies that A cuts each stratum.

Example 9. Examples with an infinite value for Ocat can be obtained easily. For space X , we take the plane \mathbb{R}^2 stratified by the orbits of the action of the rotation group $\text{SO}(2)$. Let A be the half-ray Ox . We observe that any open subset U containing the singular orbit O contains also a circle C which cannot be contracted to some point of $A \cap C$ by a stratified deformation of U . Therefore $\text{Ocat}(\mathbb{R}^2, Ox, \mathcal{F}_{\mathbb{R}^2}) = \infty$.

The next easy result will be used in the proof of Theorem D.

Proposition 10. *Let (X, A, \mathcal{F}_X) be a stratified pair. If $f: (X, \mathcal{F}_X) \rightarrow (Y, \mathcal{F}_Y)$ is a stratified map with a right \mathcal{S} -homotopical inverse g , then*

$$\text{Ocat}(Y, f(A), \mathcal{F}_Y) \leq \text{Ocat}(X, A, \mathcal{F}_X).$$

Proof. Let $U \subseteq X$ be an A -categorical open set and $H: (U, \mathcal{F}_U) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)$ be a stratified deformation of U into A . On the open set $V = g^{-1}(U)$, we define a stratified deformation $f \circ H \circ (g \times \text{id}_I): (V, \mathcal{F}_V) \times (I, \mathcal{F}_I) \rightarrow (Y, \mathcal{F}_Y)$ into $f(A)$. The inequality follows. \square

2.4. Comparison of the three invariants.

Theorem B. *For any stratified pair (X, A, \mathcal{F}_X) , we have the equality*

$$\text{Whcat}(X, A, \mathcal{F}_X) = \text{Gcat}(X, A, \mathcal{F}_X).$$

Proof. We first define a stratified map $\varepsilon_n: G_n(X, A, \mathcal{F}_X) \rightarrow T_n(X, A, \mathcal{F}_X)$ by $\varepsilon_n(\alpha_1, \dots, \alpha_{n+1}) = (\alpha_1(0), \dots, \alpha_{n+1}(0))$. The following diagram is clearly homotopy commutative:

$$\begin{array}{ccc} G_n(X, A, \mathcal{F}_X) & \xrightarrow{g_n} & (X, \mathcal{F}_X) \\ \varepsilon_n \downarrow & & \downarrow \Delta \\ T_n(X, A, \mathcal{F}_X) & \xrightarrow{t_n} & (X^{n+1}, \mathcal{F}_{X^{n+1}}) \end{array}$$

The proof is reduced to the fact that this diagram is a homotopy pull-back. For that, we need to determine the associated fibration to the diagonal Δ , which is obtained from the pull-back

$$\begin{array}{ccc} (P, \mathcal{F}_P) & \longrightarrow & (X^{n+1}, \mathcal{F}_{X^{n+1}})^I \\ \downarrow & & \downarrow \\ (X, \mathcal{F}_X) & \xrightarrow{\Delta} & (X^{n+1}, \mathcal{F}_{X^{n+1}}) \end{array}$$

By construction, (P, \mathcal{F}_P) is the space of $(n+1)$ -uples $(\alpha_1, \dots, \alpha_{n+1})$ of paths in $(X, \mathcal{F}_X)^I$, such that $\alpha_1(1) = \dots = \alpha_{n+1}(1)$. The evaluation map to 0 of these paths in X^{n+1} is a fibration equivalent to the diagonal. So if we take the pull-back of this map and of the map $t_n: T_n(X, A, \mathcal{F}_X) \rightarrow (X^{n+1}, \mathcal{F}_{X^{n+1}})$, we get the homotopy pull-back of t_n and of the diagonal. But this is exactly the Ganea space $G_n(X, A, \mathcal{F}_X)$. \square

We prove now the equivalence between the Whitehead and the open-set definitions, under some hypotheses, as in the classical case where one needs the existence of a contractible neighborhood of the base point. The corresponding notion is given by the following definition.

Definition 11. Let (X, \mathcal{F}_X) be a stratified space and let A, B be subsets of X . We say that A is a B stratified neighborhood deformation (in short B -SND) if A has some open neighborhood which is a B -categorical set.

Theorem C. Let (X, \mathcal{F}_X) be a stratified space and A be a subspace of X .

(1) If the space X is normal, then we have the inequality:

$$\text{Whcat}(X, A, \mathcal{F}_X) \leq \text{Ocat}(X, A, \mathcal{F}_X).$$

(2) If A is a B -SND, then we have the inequality:

$$\text{Ocat}(X, B, \mathcal{F}_X) \leq \text{Whcat}(X, A, \mathcal{F}_X).$$

Proof. 1) Suppose $\text{Ocat}(X, A, \mathcal{F}_X) \leq n$. There is a covering of X by open sets, U_0, \dots, U_n , and stratified deformations $H_i: (U_i, \mathcal{F}_{U_i}) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)$ into A , for $i = 0, \dots, n$. As X is normal, there exists a covering of X by open sets, W_0, \dots, W_n , such that $\overline{W}_i \subset U_i$, for $i = 0, \dots, n$. For any i , we choose an Urysohn function $\varphi_i: X \rightarrow I$ such that $\varphi_i(x) = 1$ if $x \in \overline{W}_i$ and $\varphi_i(x) = 0$ if $x \notin U_i$. We define now a continuous stratified map $\hat{H}_i: (X, \mathcal{F}_X) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)$ by

$$\hat{H}_i(x, t) = \begin{cases} H_i(x, \varphi_i(x)t) & \text{if } x \in U_i, \\ x & \text{otherwise.} \end{cases}$$

We collect these maps in a continuous stratified map $H: (X, \mathcal{F}_X) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)^{n+1}$ defined by $H(x, t) = (\hat{H}_0(x, t), \dots, \hat{H}_n(x, t))$. Observe that we have $H(x, 0) = (x, \dots, x) = \Delta(x)$.

Set $r(x) = H(x, 1)$. Since the W_i 's are a covering of X , for any point $x \in X$, there is a W_j with $x \in W_j$. By definition of \hat{H}_j , we have $\hat{H}_j(x, 1) = H_j(x, 1) \in A$. From the construction of the fat wedge, we have $r(X) \subset T_n(X, A, \mathcal{F}_X)$ and r is a lifting up to \mathcal{S} -homotopy (by the \mathcal{S} -homotopy H) of the diagonal. By definition, we get $\text{Whcat}(X, A, \mathcal{F}_X) \leq n$.

2) Suppose $\text{Whcat}(X, A, \mathcal{F}_X) \leq n$. By definition, there are a stratified map $r: (X, \mathcal{F}_X) \rightarrow T_n(X, A, \mathcal{F}_X)$ and a stratified homotopy $H: (X, \mathcal{F}_X) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)^{n+1}$ between the diagonal Δ and the composite $t_n \circ r$, see Definition 6. As A is a B -SND, there exists also an open set N , $A \subset N$, and a stratified homotopy $G: (N, \mathcal{F}_N) \times (I, \mathcal{F}_I) \rightarrow (X, \mathcal{F}_X)$ of N with $G(x, 1) \in B$.

Let $p_i: X^{n+1} \rightarrow X$ be the $(i+1)$ th projection, $0 \leq i \leq n$, we set $h_i = p_i \circ t_n \circ r$ and $U_i = h_i^{-1}(N)$. Then, since $r(X) \subset T_n(X, A, \mathcal{F}_X) = \bigcup_{i=0}^n p_i^{-1}(A)$ we have $X = \bigcup_{i=0}^n U_i$. The U_i 's are a covering of X . They are B -categorical with the homotopy $H_i: U_i \times I \rightarrow X$ defined by

$$H_i(u, t) = \begin{cases} p_i H(u, 2t) & \text{if } 0 \leq t \leq 1/2, \\ G(h_i(u), 2t - 1) & \text{if } 1/2 \leq t \leq 1. \end{cases}$$

It is routine to check that H_i is a continuous homotopy between the identity and a map with values in B . As H , p_i , G and h_i are stratified maps, H_i is stratified also and we have proved $\text{Ocat}(X, B, \mathcal{F}_X) \leq n$. \square

Let X be path-connected with the trivial stratification and let U be a categorical open set in X . For any point $u \in U$ and any $x \in X$, the subset $\{u\}$ is an $\{x\}$ -SND. But, in the general case of a stratified space, being B -SND depends on the stratification but also depends on B . This explains why we need the notion of transverse sets, introduced in the next section.

3. LUSTERNIK-SCHNIRELMANN CATEGORY OF STRATIFIED SPACES

First, we define the *transverse subsets* of a stratified space (X, \mathcal{F}_X) which play a fundamental role for the comparison of our previous invariants with the tangential LS-category of foliations.

Definition 12. Let (X, \mathcal{F}_X) be a stratified space. A subspace $A \subseteq X$ is *transverse to the stratification* if, for any stratum $S \in \mathcal{F}_X$, the set $A \cap S$ is at most countable. We denote this property by $A \pitchfork \mathcal{F}_X$.

As it is proved by W. Singhof and E. Vogt, [15], the transverse subspaces of a foliation are transverse in the sense of Definition 12. We observe now that, in Definition 18 of the tangential category of a foliation, the transverse space which receives the stratified deformation is not predetermined. We have therefore to take in account all the transverse subspaces associated to tangential deformations.

Definition 13. Let (X, \mathcal{F}_X) be a stratified space. The *open LS-category* of (X, \mathcal{F}_X) is the infimum of the integers $\text{Ocat}(X, A, \mathcal{F}_X)$, when A runs along the transverse subsets to \mathcal{F}_X :

$$\text{Ocat}(X, \mathcal{F}_X) = \text{Inf} \{ \text{Ocat}(X, A, \mathcal{F}_X) \mid A \pitchfork \mathcal{F}_X \}.$$

Similarly, we define the *Whitehead LS-category* and the *Ganea LS-category* of the stratified space (X, \mathcal{F}_X) by taking the infimum along the transverse subsets. We denote them by $\text{Gcat}(X, \mathcal{F}_X)$ and $\text{Whcat}(X, \mathcal{F}_X)$ respectively.

The lower bound gives a homotopy invariant.

Theorem D. *In the fibration category $\mathcal{S}\text{-Top}$, the integer $\text{Ocat}(-, -)$ is a homotopy invariant.*

Proof. Let $f: (X, \mathcal{F}_X) \rightarrow (X', \mathcal{F}_{X'})$ be a stratified map with homotopical inverse g in $\mathcal{S}\text{-Top}$. Let \sim (resp. \sim') be the equivalence relation generated by \mathcal{F}_X (resp. $\mathcal{F}_{X'}$). The stratum containing $x \in X$ (resp. $f(x) \in X'$) is denoted by S_x (resp. $S'_{f(x)}$). The proof is divided in several steps.

1) First, one proves that f induces a homeomorphism $\bar{f}: X/\sim \rightarrow X'/\sim'$ between the quotient spaces. If $y \in S'_{f(x)}$, there is a path $\gamma: [0, 1] \rightarrow S'_{f(x)}$ joining y and $f(x)$. Therefore $g(y)$ and $g(f(x))$ are connected by a path included in one stratum. As $g \circ f \simeq_S \text{id}_X$, the points $g(f(x))$ and x are in the same stratum. This implies $g(S'_{f(x)}) \subseteq S_x$ and $\bar{g} \circ \bar{f} = \text{id}_{X/\sim}$. With a similar argument, one has $\bar{f} \circ \bar{g} = \text{id}_{X'/\sim'}$.

2) Suppose $f(S) \subseteq S'$ where S and S' are strata of \mathcal{F}_X and $\mathcal{F}_{X'}$ respectively. Let $A \subset X$. If $f(a) \in f(A) \cap S'$, then $g(f(a)) \in S$, from part (1), and $a \in S$, because a and $g(f(a))$ are in the same stratum. This implies $f(A) \cap S' \subseteq f(A \cap S)$. As the reverse inclusion is obvious, one has $f(A \cap S) = f(A) \cap S'$.

3) Let $A \subset X$ such that $f(A)$ is not transverse to $\mathcal{F}_{X'}$, i.e. the set $f(A) \cap S'$ is uncountable for some stratum $S' \in \mathcal{F}_{X'}$. Let $S \in \mathcal{F}_X$ with $g(S') \subseteq S$. From the equality $f(A \cap S) = f(A) \cap S'$, we deduce that $A \cap S$ is uncountable. We have proved that $A \pitchfork \mathcal{F}_X$ implies $f(A) \pitchfork \mathcal{F}_{X'}$.

4) Proposition 10 implies the inequality

$$\text{Inf} \{ \text{Ocat}(X, A, \mathcal{F}_X) \mid A \pitchfork \mathcal{F}_X \} \geq \text{Inf} \{ \text{Ocat}(X', A', \mathcal{F}_{X'}) \mid A' \pitchfork \mathcal{F}_{X'} \}.$$

As the opposite inequality follows by symmetry, we have proved the equality $\text{Ocat}(X, \mathcal{F}_X) = \text{Ocat}(X', \mathcal{F}_{X'})$. \square

Remark 14. In the previous proof, we have used, in a fundamental way, the fact that the image, by a weak equivalence of $\mathcal{S}\text{-Top}$, of a transverse set of (X, \mathcal{F}_X) is a transverse set of $(X', \mathcal{F}_{X'})$. With the next example, we observe that this is not true for the subsets A of (X, \mathcal{F}_X) such that $A \cap S$ is totally disconnected for any $S \in \mathcal{F}_X$. For similar reasons, the property $A \cap S$ of topological dimension 0 does not fit also, see [12, Theorem, p. 302].

Example 15. Let $f: X = S^1 \times \mathbb{R} \rightarrow X' = S^1$ be the projection on the first factor. We put on X and X' the trivial stratification with only one stratum. Let $\varphi: S^1 \rightarrow \mathbb{R}$ be an application that is discontinuous at each point and let A be the graph of φ . We see easily that f is a weak equivalence of $\mathcal{S}\text{-Top}$, that A is totally disconnected and $f(A) = X'$ is not.

4. TANGENTIAL LS-CATEGORY OF FOLIATIONS

The *tangential LS-category* of a foliated manifold was introduced by H. Colman and the second author in [6], see also [3]. In this section we compare it with our previous definitions of stratified LS-categories; all our manifolds are C^∞ -manifolds.

Let M be a foliated manifold of class C^0 . The leaves of the foliation form a partition \mathcal{F}_M of M . Moreover the definitions of foliated continuous maps and tangential continuous homotopies correspond exactly to our definitions of stratified maps and stratified homotopies. The induced foliation on an open set of M coincides also with our induced stratification of Definition 2. In the case of a foliation of class C^∞ on a closed manifold, recall from [15, Theorem 4.1] that the use of continuous maps or of C^∞ -maps does not modify the tangential LS-category.

4.1. Definition of tangential LS-category.

Definition 16. An open set $U \subset M$ (endowed with the induced foliation \mathcal{F}_U) is *tangentially categorical* if there exists a tangential homotopy $H: U \times I \rightarrow M$ such that $H(-, 0)$ is the inclusion and $H(-, 1)$ is constant along the leaves of U , that is $H(-, 1)$ maps each leaf of \mathcal{F}_U onto some point. We call H a *tangential contraction*.

Example 17. In general a tangential contraction defined on the categorical open set U cannot be extended to the whole manifold M . For instance, let M be the punctured plane $\mathbb{R}^2 \setminus \{(0, 0)\}$, foliated by the horizontal lines $\mathbb{R} \times \{t\}$. Let L be the closed leaf $L = \{(x, 0) : x < 0\}$ and take $U = M \setminus L$. Then U can be tangentially contracted to $A = \{1\} \times \mathbb{R}$, but the contraction cannot be extended to M .

Definition 18. [3, 6] The *tangential LS-category* of the foliated manifold (M, \mathcal{F}_M) is the least integer n such that there exists a covering of M by $(n+1)$ open sets which are tangentially categorical. We denote it by $\text{cat}_{\mathcal{F}}(M)$.

4.2. Equivalence of the four invariants. This paragraph is devoted to the proof of the equivalence between the tangential LS-category of a foliation and our definitions of the LS-category of the corresponding stratified space. Let (M, \mathcal{F}_M) be a foliated manifold of class C^0 and dimension p .

Lemma 19. *Consider a tangentially categorical open set U with its tangential contraction $H: U \times I \rightarrow M$. Then $H(U, 1)$ is contained in a set $A(U)$ which is a finite or countable union of compact $(n-p)$ -submanifolds with boundary, each one transverse to \mathcal{F}_M . Moreover, for each leaf L of \mathcal{F}_M the intersection $L \cap A(U)$ is a countable set.*

Proof. The first part is [15, Lemma 1.1]. The second part follows from the fact that each compact transverse submanifold can be covered by a finite number of adapted charts. Since its dimension is the codimension of the foliation, its intersection with a leaf is a second countable manifold of dimension zero. \square

Theorem E. *If \mathcal{F}_M is a C^0 -foliation on the manifold M , we have the equality $\text{Ocat}(M, \mathcal{F}_M) = \text{cat}_{\mathcal{F}}(M)$. In the case of a C^1 foliation, the four invariants coincide:*

$$\text{Gcat}(M, \mathcal{F}_M) = \text{Whcat}(M, \mathcal{F}_M) = \text{Ocat}(M, \mathcal{F}_M) = \text{cat}_{\mathcal{F}}(M).$$

Proof. We first prove that $\text{cat}_{\mathcal{F}}(M) = \text{Ocat}(M, \mathcal{F}_M)$.

- Suppose $\text{cat}_{\mathcal{F}}(M) < \infty$. Let U_0, \dots, U_n be a covering of M by tangentially categorical open sets. Denote by $H_i: U_i \times I \rightarrow M$ a tangential contraction of U_i and by A_i the set $A(U_i)$ containing $H_i(U_i, 1)$ as in Lemma 19. Let $A = A_0 \cup \dots \cup A_n$. Then the set A is transverse to \mathcal{F}_M in the sense of Definition 12, and the open sets U_i are A -categorical (Definition 7). Hence $\text{Ocat}(M, \mathcal{F}_M) \leq n$.

- On the other hand, suppose $\text{Ocat}(M, \mathcal{F}_M) < \infty$. Since this infimum must be a minimum, there exists a transverse subset A such that $\text{Ocat}(M, \mathcal{F}_M) = \text{Ocat}(M, A, \mathcal{F}_M)$. Let U be an A -categorical open set and H be the corresponding stratified homotopy. Then for each leaf $L \in \mathcal{F}_M$, the intersection $L \cap A$ is a countable, thus totally disconnected set. Hence the image by $H(-, 1)$ of each connected component of $L \cap U$ is reduced to a point. Therefore, U is tangentially categorical, which proves that $\text{cat}_{\mathcal{F}}(M) \leq n$.

Now, Theorems B and C (part (1)) give


$$\text{Gcat}(M, \mathcal{F}_M) = \text{Whcat}(M, \mathcal{F}_M) \leq \text{Ocat}(M, \mathcal{F}_M) = \text{cat}_{\mathcal{F}}(M).$$

So we just have to prove that $\text{Ocat}(M, \mathcal{F}_M) \leq \text{Whcat}(M, \mathcal{F}_M)$.

- If $\text{Whcat}(M, \mathcal{F}_M) = n$, there exists A such that $\text{Whcat}(M, A, \mathcal{F}_M) = n$ and $A \pitchfork \mathcal{F}_M$. By Lemma 1.1 and Proposition 5.1 of [15], there is a subset B , such that $B \pitchfork \mathcal{F}_M$ and A is a B -SND. Now Theorem C implies the inequality $\text{Ocat}(M, \mathcal{F}_M) \leq \text{Whcat}(M, \mathcal{F}_M)$. \square

Example 20 (Reeb foliation). We ask if for an arbitrary foliation there exists a transverse subset A such that we have the equality $\text{Ocat}(M, \mathcal{F}_M) =$

$\text{Ocat}(M, A, \mathcal{F}_M)$ and that is an A -SND, i.e. a SND of itself. Here we construct explicitly such A for the example of the Reeb foliation on T^2 , as described in [6, Section 6.2]. Consider the following representation of the torus where the upper and lower arcs are identified. At the bottom, the left and right segments have also to be identified. The strata are the horizontal lines.



In each of these two pictures, a tangential categorical open set is formed of the colored and hatched areas together with their symmetric relatively to $\alpha \cup \beta$. Denote it by U . It is deformed into a transverse set A which is the union of the curves α, β, γ . In the second picture A lies inside U , showing that A is an A -SND. The tangential deformation of U into A is defined as follows:

- the dark gray zone of U is deformed on the upper part α of the vertical line;
- the clear gray zone is deformed on the lower part β of the vertical line;
- the hatched zone is deformed on the curve γ .

REFERENCES

[1] Hans Joachim Baues. *Algebraic homotopy*, volume 15 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1989.

[2] Mónica Clapp and Dieter Puppe. Invariants of the Lusternik-Schnirelmann type and the topology of critical sets. *Trans. Amer. Math. Soc.*, 298:603–620, 1986.

[3] Hellen Colman. *Categoría L.S en foliaciones*. Publ. Dep. Geom. y Topología, Vol. 93, Santiago de Compostela, 1998.

[4] Hellen Colman and Steven Hurder. Tangential LS category and cohomology for foliations. In *Lusternik-Schnirelmann category and related topics (South Hadley, MA,*

- 2001), volume 316 of *Contemp. Math.*, pages 41–64. Amer. Math. Soc., Providence, RI, 2002.
- [5] Hellen Colman and Enrique Macias-Virgós. Transverse Lusternik-Schnirelmann category of foliated manifolds. *Topology*, 40(2):419–430, 2001.
 - [6] Hellen Colman and Enrique Macias-Virgós. Tangential Lusternik-Schnirelmann category of foliations. *J. London Math. Soc. (2)*, 65(3):745–756, 2002.
 - [7] Octavian Cornea, Gregory Lupton, John Oprea, and Daniel Tanré. *Lusternik-Schnirelmann category*, volume 103 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
 - [8] Jean-Paul Doeraene. L.S.-category in a model category. *J. Pure Appl. Algebra*, 84(3):215–261, 1993.
 - [9] Albrecht Dold. Partitions of unity in the theory of fibrations. *Annals of Math.*, 78(2):223–255, 1963.
 - [10] Agnese Fasso Velenik. Relative homotopy invariants of the type of the Lusternik-Schnirelmann category. Thesis, Freie Universität Berlin, 2003.
 - [11] Steven Hurder. Foliations geometry/topology problem set. Survey, prepared for the conference Geometry and Foliations held at Ryukoku University in Kyoto, Japan, September 10-19, 2003. Available at <http://www.math.uic.edu/~hurder>.
 - [12] Kasimierz Kuratowski. *Topology. Vol. I*. New edition, revised and augmented. Translated from the French by J. Jaworowski. Academic Press, New York, 1966.
 - [13] Martin Majewski. Rational homotopical models and uniqueness. *Mem. Amer. Math. Soc.*, 143(682):xviii+149, 2000.
 - [14] Daniel G. Quillen. *Homotopical algebra*. Lecture Notes in Mathematics, No. 43. Springer-Verlag, Berlin, 1967.
 - [15] Wilhelm Singhof and Elmar Vogt. Tangential category of foliations. *Topology*, 42(3):603–627, 2003.
 - [16] Arne Strøm. The homotopy category is a homotopy category. *Arch. Math. (Basel)*, 23:435–441, 1972.

DÉPARTEMENT DE MATHÉMATIQUES, UMR-CNRS 8524, UNIVERSITÉ DE LILLE 1,
59655 VILLENEUVE D’ASCQ CEDEX, FRANCE

E-mail address: Jean-Paul.Doeraene@univ-lille1.fr

DEPARTAMENTO DE XEOMETRIA E TOPOLOXIA, FACULTADE DE MATEMÁTICAS, UNI-
VERSIDADE DE SANTIAGO DE COMPOSTELA, 15706 GALICIA, SPAIN

E-mail address: quique.macias@usc.es

DÉPARTEMENT DE MATHÉMATIQUES, UMR-CNRS 8524, UNIVERSITÉ DE LILLE 1,
59655 VILLENEUVE D’ASCQ CEDEX, FRANCE

E-mail address: Daniel.Tanre@univ-lille1.fr